arXiv:hep-ph/0403136v1 12 Mar 2004

Covariant tensor formalism for partial wave analyses of ψ decays into $\gamma B\bar{B}$

Sayipjamal Dulat[b,c], Bo-Chao Liu[b] , Bing-Song Zou[a,b,], and Ji-Min Wu[a,b,]

a) CCAST (World Laboratory), P.O. Box 8730, Beijing 100080

b) Institute of High Energy Physics, CAS, P.O. Box 918(4), Beijing, 100039, China

c)Department of Physics, Xinjiang University, Urumqi, 830046, China

Abstract

Recently BES2 collaboration observed an enhancement near the $p\bar{p}$ invariant mass spectrum. Using the covariant tensor formalism, here we provide theoretical formulae for the partial wave analysis (PWA) of the ψ radiative decay channels $\psi \to \gamma p\bar{p}, \gamma\Lambda\bar{\Lambda}, \gamma\Sigma\bar{\Sigma}, \gamma\Xi\bar{\Xi}$. By performing the Monte Carlo simulation, we give the angular distributions for the photon and proton in the process of $J/\psi \to \gamma p\bar{p}$, which may serve as a useful reference for the future PWA on these channels.

1 Introduction

J/ψ and ψ' radiative decay to $B\bar{B}$ (baryon and antibaryon pair) is a good channel to study the possible bound or resonant states of the $B\bar{B}$ system. Abundant J/ψ and ψ' events have been collected at the Beijing Electron Positron Collider (BEPC). More data will be collected at upgraded BEPC and CLEO-C. Based on the analysis of the 58 million J/ψ events accumulated by the BES2 detector at the BEPC, recently BES2 reported [5] that they observed a strong, narrow enhancement near the threshold in the invariant mass spectrum of $p\bar{p}$ (proton - antiproton) pairs from $J/\psi \to \gamma p\bar{p}$ radiative decays. The structure has the properties consistent with either an $S(0^{-+})$- or $P(0^{++})$-wave Breit-Wigner resonance function. In the S-wave case, the peak mass is blow $2M_p$ around $M = 1859MeV$ with a total width $\Gamma < 30MeV$.In order to get more useful information about properties of the resonances such as their J^{PC} quantum numbers, mass, width, production and decay rates, etc., partial wave analyses (PWA) are necessary. PWA is an effective method for analysing the experimental data of hadron spectrum. There are two types of PWA: one is based on the covariant tensor (also named Rarita-Schwinger) formalism[6] and the other is based on the helicity formalism[7]. Ref.[8] showed the connection between the covariant tensor formalism and the covariant helicity one. Ref.[9] provided PWA formulae in a covariant tensor formalism for ψ decays to mesons, which have been used for a number of channels already published by BES[10, 11, 12, 13, 14, 15]

and are going to be used for more channels. A similar approach has been used in analyzing other reactions[16, 17, 18]. Ref.[19] provided explicit formulae for the angular distribution of photon of the ψ radiative decays in the covariant tensor formalism, and also discussed helicity formalism of the angular distribution of the ψ radiative decays to two pseudoscalar mesons, and its relation to the covariant tensor formalism. Now we extend the covariant tensor formalism to the $\psi \to \gamma B\bar{B}$ with B represents baryons p, Λ, Σ, Ξ etc, and derive the decay amplitudes for various intermediate resonant states in the framework of the relativistic covariant tensor formalism.

In this paper we study the phenomenological spin parity determination of resonances. The plan of this article is as follows: in section 2, we present the necessary tools for the calculation of the tensor amplitudes, within a covariant tensor formalism. This will allow us to derive covariant amplitudes for all possible processes. In section 3, we present covariant tensor formalism for ψ radiative decays to baryon antibaryon pairs. Since covariance is a general requirement of any decay amplitude, all possible amplitudes are written in terms of covariant tensor form. All amplitudes include a complex coupling constant and Blatt-Weisskopf centrifugal barriers where necessary. In section 3, we provide the angular distribution of the photon and proton. The conclusions are given in section 5.

2 Prescriptions for the construction of covariant tensor amplitudes

In this section we present the necessary tools for the construction of covariant tensor amplitudes. The partial wave amplitudes $U_i^{\mu\nu\alpha}$ in the covariant Rarita-Schwinger tensor formalism can be constructed by using pure orbital angular momentum covariant tensors $\tilde{t}^{(L_{bc})}_{\mu_1\cdots\mu_{L_{bc}}}$ and covariant spin wave functions $\phi_{\mu_1\cdots\mu_s}$ together with metric tensor $g^{\mu\nu}$, totally antisymmetric Levi-Civita tensor $\epsilon_{\mu\nu\lambda\sigma}$ and momenta of parent particles. For a process $a \to bc$, if there exists a relative orbital angualar momentum $\mathbf{L}_{bc}$ between the particle a and b, then the pure orbital angular momentum $\mathbf{L}_{bc}$ state can be represented by covariant tensor wave functions $\tilde{t}^{(L_{bc})}_{\mu_1\cdots\mu_{L_{bc}}}$[7] which is built out of relative momentum. Thus here we give only covariant tensors that correspond to the pure S-, P-, D-, and F-wave orbital angular momenta:

$$\tilde{t}^{(0)} = 1, \tag{1}$$

$$\tilde{t}^{(1)}_{\mu} = \tilde{g}_{\mu\nu}(p_a) r^{\nu} B_1(Q_{abc}) \equiv \tilde{r}_{\mu} B_1(Q_{abc}), \tag{2}$$

$$\tilde{t}^{(2)}_{\mu\nu} = [\tilde{r}_{\mu}\tilde{r}_{\nu} - \frac{1}{3}(\tilde{r}\cdot\tilde{r})\tilde{g}_{\mu\nu}(p_a)] B_2(Q_{abc}), \tag{3}$$

$$\tilde{t}^{(3)}_{\mu\nu\lambda} = [\tilde{r}_{\mu}\tilde{r}_{\nu}\tilde{r}_{\lambda} - \frac{1}{5}(\tilde{r}\cdot\tilde{r})(\tilde{g}_{\mu\nu}(p_a)\tilde{r}_{\lambda} + \tilde{g}_{\nu\lambda}(p_a)\tilde{r}_{\mu} + \tilde{g}_{\lambda\mu}(p_a)\tilde{r}_{\nu})] B_3(Q_{abc}), \tag{4}$$

...

where $r = p_b - p_c$ is the relative four momentum of the two decay products in the parent particle rest frame; $(\tilde{r} \cdot \tilde{r}) = -\mathbf{r}^2$. and

$$\tilde{g}_{\mu\nu}(p_a) = g_{\mu\nu} - \frac{p_{a\mu}p_{a\nu}}{p_a^2}; \tag{5}$$

$B_{L_{bc}}(Q_{abc})$ is a Blatt-Weisskopf barrier factor[7, 21], here Q_{abc} is the magnitude of $\mathbf{p_b}$ or $\mathbf{p_c}$ in the rest system of a,

$$Q_{abc}^2 = \frac{(s_a + s_b - s_c)^2}{4s_a} - s_b \tag{6}$$

with $s_a = E_a^2 - \mathbf{p}_a^2$.

The spin-1 and spin-2 particles wave functions $\phi_\mu(p_a, m)$ and $\phi_{\mu\nu}(p_a, m)$ satisfy the following conditions

$$\begin{aligned} &p_a^\mu \phi_\mu(p_a, m) = 0, \qquad \phi_\mu(p_a, m)\phi^{*\mu}(p_a, m) = -\delta_{mm'} \\ &\sum_m \phi_\mu(p_a, m)\phi_\nu^*(p_a, m) = -g_{\mu\nu} + \frac{p_{a\mu}p_{a\nu}}{p_a^2} \equiv -\tilde{g}_{\mu\nu}(p_a), \end{aligned} \tag{7}$$

and

$$\begin{aligned} &p_a^\mu \phi_{\mu\nu}(p_a, m) = 0, \qquad \phi_{\mu\nu} = \phi_{\nu\mu}, \qquad g^{\mu\nu}\phi_{\mu\nu} = 0, \qquad \phi_{\mu\nu}(m)\phi^{*\mu\nu}(m') = \delta_{mm'} \\ &P^{(2)}_{\mu\nu\mu'\nu'}(p_a) = \sum_m \phi_{\mu\nu}(p_a, m)\phi^*_{\mu'\nu'}(p_a, m) = \frac{1}{2}(\tilde{g}_{\mu\mu'}\tilde{g}_{\nu\nu'} + \tilde{g}_{\mu\nu'}\tilde{g}_{\nu\mu'}) - \frac{1}{3}\tilde{g}_{\mu\nu}\tilde{g}_{\mu'\nu'}. \end{aligned} \tag{8}$$

Note that for a given decay process $a \to bc$, the total angular momentum should be conserved, which means

$$\mathbf{J}_a = \mathbf{S}_{bc} + \mathbf{L}_{bc}, \tag{9}$$

where

$$\mathbf{S}_{bc} = \mathbf{S}_b + \mathbf{S}_c. \tag{10}$$

In addition parity should also be conserved, which means

$$\eta_a = \eta_b\eta_c(-1)^{L_{bc}}, \tag{11}$$

where η_a, η_b, and η_c are the intrinsic parities of particles a, b, and c, respectively. From this relation, one knows whether L_{bc} should be even or odd. Then from Eq. (9) one can find out how many different $L_{bc} - S_{bc}$ combinations, which determine the number of independent couplings. Also note that in the construction of the covariant tensor amplitude, for $S_{bc} - L_{bc} - J_a$ coupling, if $S_{bc} + L_{bc} + J_a$ is an odd number, then $\epsilon_{\mu\nu\lambda\sigma}p_a^\sigma$ with p_a the momentum of the parent particle is needed; otherwise it is not needed.

3 Covariant tensor formalism for ψ decay into $\gamma B\bar{B}$

The general form of the decay $\psi \to \gamma X \to \gamma p\bar{p}$ amplitude can be written as follows by using the polarization four-vectors of the initial and final states,

$$A^{(s)} = \psi_\mu(m_J)e^*_\nu(m_\gamma)\psi_{\alpha_s}(p_b,S_b;p_c,S_c)A^{\mu\nu\alpha_s} = \psi_\mu(m_J)e^*_\nu(m_\gamma)\psi_{\alpha_s}(p_b,S_b;p_c,S_c)\sum_i \Lambda_i U_i^{\mu\nu\alpha_s}. \tag{12}$$

where $\psi_\mu(m_J)$ is the polarization four vector of the ψ with spin projection of m_J; $e_\nu(m_\gamma)$ is the polarization four vector of the photon with spin projections of m_γ; $U_i^{\mu\nu\alpha_s}$ is the i-th partial wave amplitude with coupling strength determined by a complex parameter Λ_i. The spin-1 polarization vector $\psi_\mu(m_J)$ for ψ with four momentum p_μ satisfies

$$\sum_{m_J=1}^{3}\psi_\mu(m_J)\psi^*_\nu(m_J) = -g_{\mu\nu} + \frac{p_\mu p_\nu}{p^2} \equiv -\tilde{g}_{\mu\nu}(p), \tag{13}$$

with $p^\mu\psi_\mu = 0$. Here the Minkowsky metric tensor has the form

$$g_{\mu\nu} = diag(1,-1,-1,-1).$$

For ψ production from e^+e^- annihilation, the electrons are highly relativistic, with the result that $J_z = \pm 1$ for the ψ spin projection taking the beam direction as the z-axis. This limits m_J to 1 and 2, i.e. components along x and y. Then one has the following relation

$$\sum_{m_J=1}^{2}\psi_\mu(m_J)\psi^*_{\mu'}(m_J) = \delta_{\mu\mu'}(\delta_{\mu 1}+\delta_{\mu 2}). \tag{14}$$

For the photon polarization four vector, there is the usual Lorentz orthogonality conditions. Namely, the polarization four vector $e_\nu(m_\gamma)$ of the photon with momenta q satisfies

$$q^\nu e_\nu(m_\gamma) = 0, \tag{15}$$

which states that spin-1 wave function is orthogonal to its own momentum. The above relation is the same as for a massive vector meson. However, for the photon, there is an additional gauge invariance condition. Here we assume the Coulomb gauge in the ψ rest system, *i.e.*, $p^\nu e_\nu = 0$. Then we have [20]

$$\sum_{m_\gamma} e^*_\mu(m_\gamma)e_\nu(m_\gamma) = -g_{\mu\nu} + \frac{q_\mu K_\nu + K_\mu q_\nu}{q\cdot K} - \frac{K\cdot K}{(q\cdot K)^2}q_\mu q_\nu \equiv -g^{(\perp\perp)}_{\mu\nu} \tag{16}$$

with $K = p - q$ and $K^\nu e_\nu = 0$. For $X \to p\bar{p}$, the total spin of $p\bar{p}$ system can be either 0 or 1. These two states can be represented by ψ and ψ_α [22]. where

$$\psi = \bar{u}(p_b,S_b)\gamma_5 v(p_c,S_c), \qquad if \quad s=0, \tag{17}$$

$$\psi_\alpha = \bar{u}(p_b,S_b)(\gamma_\alpha - \frac{r_\alpha}{m_X+2m})v(p_c,S_c), \qquad if \quad s=1. \tag{18}$$

One can see that both ψ and ψ_α have no dependence on the direction of the momentum $\hat{\mathbf{p}}$, hence correspond to pure spin states with the total spin of 0 and 1, respectively. Where p_b, p_c, and S_b, S_c are momenta and spin of the proton antiproton pairs, respectively. m_X and m are the masses of X and p, $\bar{p}$, respectively; $u(p_b, S_b)$ and $v(p_c, S_c)$ are the standard Dirac spinor. If we sum over the polarization, we have the two projection operators:

$$\begin{aligned}\sum_{S_b} u_\alpha(p_b, S_b)\bar{u}_\beta(p_b, S_b) &= \Big(\frac{\not{p}_b + m}{2m}\Big)_{\alpha\beta} \\ \sum_{S_c} v_\alpha(p_c, S_c)\bar{v}_\beta(p_c, S_c) &= \Big(\frac{\not{p}_c - m}{2m}\Big)_{\alpha\beta}\end{aligned} \tag{19}$$

To compute the differential cross section, we need an expression for $|A|^2$. Note that the square modulus of the decay amplitude, which gives the decay probability of a certain configuration should be independent of any particular frame, that is, a Lorentz scalar. Thus by using the Eqs. (14) and (16), the differential cross section for the radiative decay to an 3-body final state is:

$$\begin{aligned}\frac{d\sigma^{(s)}}{d\Phi_3} &= \frac{1}{2}\sum_{S_b,S_c}\sum_{m_J=1}^{2}\sum_{m_\gamma=1}^{2}|\psi_\mu(m_J)e^*_\nu(m_\gamma)\psi_{\alpha_s}(p_b, S_b; p_c, S_c)A^{\mu\nu\alpha_s}|^2 \\ &= -\frac{1}{2}\sum_{S_b,S_c}\sum_{\mu=1}^{2}A^{\mu\nu\alpha_s}g^{(\perp\perp)}_{\nu\nu'}A^{*\mu\nu'\alpha'_s}\psi^*_{\alpha_s}\psi_{\alpha'_s} \\ &= -\frac{1}{2}\sum_{i,j}\Lambda_i\Lambda^*_j\sum_{\mu=1}^{2}U^{\mu\nu\alpha_s}_i g^{(\perp\perp)}_{\nu\nu'}U^{*\mu\nu'\alpha'_s}_j\sum_{S_b,S_c}\psi^*_{\alpha_s}\psi_{\alpha'_s} \\ &\equiv \sum_{i,j}P_{ij}\cdot F^{(s)}_{ij}\end{aligned} \tag{20}$$

where

$$\begin{aligned}P_{ij} &= P^*_{ji} &&= \Lambda_i\Lambda^*_j, \\ F^{(s)}_{ij} &= F^{*(s)}_{ji} &&= -\frac{1}{2}\sum_{\mu=1}^{2}U^{\mu\nu\alpha_s}_i g^{(\perp\perp)}_{\nu\nu'}U^{*\mu\nu'\alpha'_s}_j\sum_{S_b,S_c}\psi^*_{\alpha_s}\psi_{\alpha'_s}.\end{aligned} \tag{21}$$

$d\Phi_3$ is the standard Lorentz invariant 3-body phase space given by

$$d\Phi_3(p; q, p_b, p_c) = \delta^4(p - q - p_b - p_c)\frac{d^3\mathbf{q}}{(2\pi)^3 2E_\gamma}\frac{m^2 d^3\mathbf{p}_b d^3\mathbf{p}_c}{(2\pi)^3 E_b (2\pi)^3 E_c}. \tag{22}$$

$$F^{(0)}_{ij} = F^{*(0)}_{ji} = -\frac{1}{2}\sum_{\mu=1}^{2}U^{\mu\nu}_i g^{(\perp\perp)}_{\nu\nu'}U^{*\mu\nu'}_j\sum_{S_b,S_c}\psi^*\psi$$

$$
\begin{aligned}
&= \frac{1}{2}\sum_{\mu=1}^{2} U_i^{\mu\nu} g_{\nu\nu'}^{(\perp\perp)} U_j^{*\mu\nu'} Tr\Big(\frac{\not{p}_b + m}{2m}\,\gamma_5\,\frac{\not{p}_c - m}{2m}\,\gamma_5 \Big) \\
&= -\frac{m_X^2}{4m^2}\sum_{\mu=1}^{2} U_i^{\mu\nu} g_{\nu\nu'}^{(\perp\perp)} U_j^{*\mu\nu'}. \qquad (23)
\end{aligned}
$$

The spin sums can be performed using the completeness relations from Eq. (19):

$$
\begin{aligned}
F_{ij}^{(1)} &= F_{ji}^{*(1)} = -\frac{1}{2}\sum_{\mu=1}^{2} U_i^{\mu\nu\alpha} g_{\nu\nu'}^{(\perp\perp)} U_j^{*\mu\nu'\alpha'} \sum_{S_b,S_c} \psi_\alpha^* \psi_{\alpha'} \\
&= -\frac{1}{2}\sum_{\mu=1}^{2} U_i^{\mu\nu\alpha} g_{\nu\nu'}^{(\perp\perp)} U_j^{*\mu\nu'\alpha'} \Big[Tr\Big(\frac{\not{p}_b + m}{2m}\,\gamma_\alpha\,\frac{\not{p}_c - m}{2m}\,\gamma_{\alpha'} \Big) \\
&- \frac{r_\alpha}{m_X + 2m} Tr\Big(\frac{\not{p}_b + m}{2m}\,\frac{\not{p}_c - m}{2m}\,\gamma_{\alpha'} \Big) - \frac{r_{\alpha'}}{m_X + 2m} Tr\Big(\frac{\not{p}_b + m}{2m}\,\gamma_\alpha\,\frac{\not{p}_c - m}{2m} \Big) \\
&+ \frac{r_\alpha r_{\alpha'}}{(m_X + 2m)^2} Tr\Big(\frac{\not{p}_b + m}{2m}\,\frac{\not{p}_c - m}{2m} \Big) \Big] \\
&= -\frac{1}{4m^2}\sum_{\mu=1}^{2} U_i^{\mu\nu\alpha} g_{\nu\nu'}^{(\perp\perp)} U_j^{*\mu\nu'\alpha'} \Big[p_{b\alpha} p_{b\alpha'} + p_{c\alpha} p_{c\alpha'} + p_{b\alpha} p_{c\alpha'} + p_{b\alpha'} p_{c\alpha} - m_X^2 g_{\alpha\alpha'} \Big]. \qquad (24)
\end{aligned}
$$

where we have used traces of γ matrics,

$$
\begin{aligned}
&tr(1) = 4, \qquad tr(\gamma^5) = 0, \qquad tr(\gamma^\mu\gamma^\nu) = 4g^{\mu\nu} \\
&tr(any\ \ odd\ \ \sharp\ \ \gamma' s) = 0, \qquad tr(\gamma^\mu\gamma^\nu\gamma^5) = 0. \qquad (25)
\end{aligned}
$$

Amplitudes for the radiative decay $\psi \to \gamma p\bar{p}$

We consider the decay of a ψ state into two steps: $\psi \to \gamma X$ with $X \to p\bar{p}$. The possible J^{PC} for X are $0^{++}, 0^{-+}$, 1^{++}, 2^{++}, 2^{-+}, etc. For $\psi \to \gamma X$, we choose two independent momenta p for ψ and q for the photon to be contracted with spin wave functions. We denote the four momentum of X by K. The tensor describing the first and second steps will be denoted by $\tilde{T}^{(L)}_{\mu_1\cdots\mu_L}$ and $\tilde{t}^{(l)}_{\mu_1\cdots\mu_l}$, respectively.

For $\psi \to \gamma 0^{++} \to \gamma p\bar{p}$, there is one independent covariant tensor amplitude:

$$
U^{\mu\nu\alpha} = g^{\mu\nu}\tilde{t}^{(1)\alpha}. \qquad (26)
$$

For $\psi \to \gamma 0^{-+} \to \gamma p\bar{p}$, there is one independent covariant tensor amplitude:

$$
U^{\mu\nu} = \epsilon^{\mu\nu\lambda\sigma} p_\lambda q_\sigma B_1(Q_{\psi\gamma X}). \qquad (27)
$$

For $\psi \to \gamma 1^{++} \to \gamma p\bar{p}$, there are two independent covariant tensor amplitudes:

$$U_1^{\mu\nu\alpha} = \epsilon^{\mu\nu\lambda\sigma} p_\lambda \epsilon^{\alpha\beta\rho}{}_\sigma K_\rho \tilde{t}_\beta^{(1)}, \quad (28)$$

$$U_2^{\mu\nu\alpha} = \epsilon^{\nu\lambda\sigma\gamma} p_\lambda q^\mu q_\gamma \epsilon^{\alpha\beta\rho}{}_\sigma K_\rho \tilde{t}_\beta^{(1)} B_2(Q_{\psi\gamma X}). \quad (29)$$

For $\psi \to \gamma 1^{-+}$, the exotic 1^{-+} meson cannot decay into $p\bar{p}$.
For $\psi \to \gamma 2^{++} \to p\bar{p}$, there are three independent covariant tensor amplitudes:

$$U_1^{\mu\nu\alpha} = P^{(2)\mu\nu\alpha\beta}(K) \tilde{t}_\beta^{(1)}, \quad (30)$$

$$U_2^{\mu\nu\alpha} = g^{\mu\nu} P^{(2)\alpha\beta\lambda\sigma} p_\beta p_\lambda \tilde{t}_\sigma^{(1)} B_2(Q_{\psi\gamma X}), \quad (31)$$

$$U_3^{\mu\nu\alpha} = P^{(2)\nu\alpha\beta\lambda} q^\mu p_\beta \tilde{t}_\lambda^{(1)} B_2(Q_{\psi\gamma X}). \quad (32)$$

For 2^{++} decaying to $p\bar{p}$, the orbital angular momentum between the proton and antiproton l could be 1 and 3; but we ignore $l = 3$ contribution because of the strong centrifugal barrier.

For $\psi \to \gamma 2^{-+} \to p\bar{p}$, the possible partial wave amplitudes are the following:

$$U_1^{\mu\nu} = \epsilon^{\mu\nu\lambda\sigma} p_\lambda q^\gamma \tilde{t}_{\gamma\sigma}^{(2)} B_1(Q_{\psi\gamma X}), \quad (33)$$

$$U_2^{\mu\nu} = \epsilon^{\mu\nu\lambda\sigma} p_\lambda q_\sigma p_\gamma p_\delta \tilde{t}^{(2)\gamma\delta} B_3(Q_{\psi\gamma X}), \quad (34)$$


$$U_3^{\mu\nu} = \epsilon^{\nu\gamma\lambda\sigma} p_\lambda q_\sigma q^\mu p^\delta \tilde{t}_{\gamma\delta}^{(2)} B_3(Q_{\psi\gamma X}). \quad (35)$$

It is worth to mention here that the above partial wave amplitudes for the process $J/\psi \to \gamma p\bar{p}$ are applicable to the processes $J/\psi \to \gamma\Lambda\bar{\Lambda}, \gamma\Sigma\bar{\Sigma}$, and $\gamma\Xi\bar{\Xi}$ as well.

4 Monte Carlo simulation for $J/\psi \to \gamma p\bar{p}$


We perform Monte Carlo simulation for the decay process $J/\psi \to \gamma X \to \gamma p\bar{p}$, with the J^{PC} of X to be 0^{++}, 0^{-+}, 1^{++}, 2^{++}, 2^{-+}. The predicted angular distributions for various J^{PC} intermediate resonances could serve as a useful reference for people performing partial wave analysis of $\psi \to \gamma B\bar{B}$ channels.

BES2 reported observation of a narrow enhancement near $2m_p$ in the invariant mass spectrum of $p\bar{p}$ pairs from radiative $J/\psi \to \gamma p\bar{p}$ decays. The peak has properties consistent with either a $J^{PC} = 0^{-+}$ or 0^{++}. We simulate these two processes with the Breit-Wigner mass and width of X in Ref.[5]. The angular distributions of the photon and proton are shown in Fig.1, while the invariant mass of $p\bar{p}$ and momentum distribution of the proton are shown in Fig.2. The two fits of 0^{-+} and 0^{++} are indeed hardly distinguishable from these distributions. However, if the narrow structure is really due to a narrow 0^{-+} or 0^{++} resonance, it can be distinguished by its other decay modes. While a 0^{-+} resonance can



$\cos\theta_\gamma$ $\cos\theta_p$

Figure 1: Angular distributions of the photon and proton in the process $J/\psi \to \gamma 0^{\pm +} \to \gamma p\bar{p}$.



$\mathrm{P}_p(GeV)$ $\mathrm{M}_{p\bar{p}}(GeV)$

Figure 2: Momentum distributions of proton and invariant mass of $p\bar{p}$ in the process $J/\psi \to \gamma 0^{\pm +} \to \gamma p\bar{p}$. The dashed curve is the case of 0^{++};the solid curve is the case of 0^{-+}.

decay into $\eta\pi\pi$, $K\bar{K}\pi$, a 0^{++} resonance cannot. On the other hand, while a 0^{++} resonance can decay into two pseudoscalar mesons, such as $\pi\pi$, $K\bar{K}$, a 0^{-+} resonance cannot. Both 0^{-+} and 0^{++} resonances can decay into 4π and $\pi\pi K\bar{K}$ channels. Previous data on these channels [10, 11, 12, 13, 14, 15] have not seen such narrow structure around $2m_p$. This gives support to the explanation of $p\bar{p}$ 0^{-+} final state interaction [23, 24] for the observed narrow peak structure in $p\bar{p}$ channel only.

The processes with other J^{PC} intermediate X resonances are simulated by assuming the mass of X at 2.15 GeV with a width of 0.15 GeV. The angular distribution of photon and proton are shown in Figs.3-5. The θ_γ and θ_p are given in J/ψ rest frame and X rest frame, respectively. The θ_γ angular distributions coincide with general analytic formulae given in Ref.[19] as it should be. One can see that various partial wave amplitudes give different angular distributions. By fitting the theoretical differential cross section given by Eq.(20) with parameters Λ_i to the data, one can get the magnitudes of each partial wave contribution.

5 Conclusion

In this paper we provided a theoretical formalism and a Monte Carlo study of the partial wave analysis for the radiative decay $J/\psi \to \gamma p\bar{p}$, which are also applicable to the processes $J/\psi \to \gamma\Lambda\bar{\Lambda}, \gamma\Sigma\bar{\Sigma}$ and $\gamma\Xi\bar{\Xi}$. We have constructed all possible covariant tensor amplitudes for intermediate resonant states of $J \leq 2$. For intermediate resonant states of $J \geq 3$, the production vertices need $L \geq 2$ and are expected to be suppressed [9]. The formulae here can be directly used to perform partial wave analysis of forthcoming high statistics data from CLEO-c and BES-III on these channels to extract useful information on the baryon-antibaryon interactions.

Acknowledgements

S. Dulat would like to thank Prof. Kaoru Hagiwara, who has given productive comments. Both the presentation at KEK and content of this paper has greatly benefited from his insightful comments. The work is partly supported by CAS Knowledge Innovation Project (KJCX2-SW-N02) and the National Natural Science Foundation of China under Grant Nos.10225525,10055003, 90103012, 10265003.

References

[1] C. Edwards, PhD thesis, Cal. Tech. Preprint CALT-68-1165(1985).




Figure 3: Angular distributions of the photon and proton for independent amplitudes of the process $J/\psi \to \gamma 1^{++} \to \gamma p\bar{p}$. The solid curve and dashed curve correspond to amplitudes (28) and (29), respectively.




Figure 4: Angular distributions of the photon and proton given by independent amplitudes Eq.(30) (stars), Eq.(31) (solid lines),and Eq.(32) (dashed lines) of the process $J/\psi \to \gamma 2^{++} \to \gamma p\bar{p}$.




Figure 5: Angular distributions of the photon and proton given by independent amplitudes Eq.(33) (stars), Eq.(34) (solid lines),and Eq.(35) (dashed lines) of the process $J/\psi \to \gamma 2^{-+} \to \gamma p\bar{p}$.

[2] D. Coffman et al., Phys. Rev. D41 (1990)1410.

[3] J. E. Augustin et al., Orsay preprint LAL/85-27(1985).

[4] BES Collaboration, Nucl. Phys. A675 (2000) 337 - 340.

[5] BES Collaboration, Phys. Rev. Lett. 91: 022001, 2003; hep-ex/0303006.

[6] W.Rarita, J.Schwinger, Phys. Rev. 60, 61 (1941);

[7] S.U.Chung, Phys. Rev. D48, 1225 (1993); S.U.Chung, Phys. Rev. D57,431 (1998).

[8] V. Filippini, A. Fontana, and A. Rotondi, Phys. Rev. D51, 2247 (1995).

[9] B. S. Zou and D. V. Bugg, Eur. Phys. J. A16 (2003) 537.

[10] BES Collaboration, Phys. Lett. B440 (1998) 217.

[11] BES Collaboration, Phys. Lett. B446 (1999) 356.

[12] BES Collaboration, Phys. Lett. B472 (2000) 200.

[13] BES Collaboration, Phys. Lett. B472 (2000) 208.

[14] BES Collaboration, Phys. Lett. B476 (2000) 25.

[15] BES Collaboration, Phys. Rev. D68 (2003) 052003.

[16] V.V.Anisovich, D.V.Bugg, A.V.Sarantsev, Nucl. Phys. A537 (1992) 501;
V.V.Anisovich et al., Nucl. Phys. A544 (1992) 747.

[17] A.V.Anisovich et al., Phys. Lett. B452 (1999) 173; Phys. Lett. B452 (1999) 180; Phys. Lett. B452 (1999) 187; Phys. Lett. B468 (1999) 304; Phys. Lett. B468 (1999) 309; Phys. Lett. B500 (2001) 222; Phys. Lett. B507 (2001) 23.

[18] A.V.Anisovich, V.A.Sadovnikova, Euro. Phys. J. A2 (1998) 199; A.V.Anisovich et al., J. Phys. G28 (2002) 15.

[19] Sayipjamal Dulat, Bing-Song Zou, and Ji-Min Wu, HEP& NP, Vol. 28 No.4 (2004) in press .

[20] W.Greiner et al., Quantum Chromodynamics, Springer 2002, p.242.

E.Behtends and C.Fronsdal, Phys. Rev. 106, 345 (1957).

[21] F. von Hippel and C.Quigg, Phys. Rev. D5, 624 (1972).

[22] B. S. Zou and F. Hussain, Phys. Rev. C67 015204, (2003).

[23] B. S. Zou and H. C. Chiang, Phys. Rev. D69, 034004 (2004).

[24] B. Kerbikov, A. Stavinsky, V. Fedotov, hep-ph/0402054.